

Pediatric Stroke can be frightening... Pediatric Stroke can be frightening...

1 in 4,000 babies will suffer a stroke before their first birthday, and many times before birth.

Some common signs of Pediatric Stroke:

- *Use of only one hand when playing or eating
- *Dragging one foot or walking on toes with one foot
- *Weakness on one side of the body

If you see any of these signs,
contact your doctor or your state's
early intervention program for assistance.

For more information: www.chasa.org

1 in 4,000 babies will suffer a stroke before their first birthday, and many times before birth.

Some common signs of Pediatric Stroke:

- *Use of only one hand when playing or eating
- *Dragging one foot or walking on toes with one foot
- *Weakness on one side of the body

If you see any of these signs,
contact your doctor or your state's
early intervention program for assistance.

For more information: www.chasa.org

Pediatric Stroke can be frightening... Pediatric Stroke can be frightening...

1 in 4,000 babies will suffer a stroke before their first birthday, and many times before birth.

Some common signs of Pediatric Stroke:

- *Use of only one hand when playing or eating
- *Dragging one foot or walking on toes with one foot
- *Weakness on one side of the body

If you see any of these signs,
contact your doctor or your state's
early intervention program for assistance.

For more information: www.chasa.org

1 in 4,000 babies will suffer a stroke before their first birthday, and many times before birth.

Some common signs of Pediatric Stroke:

- *Use of only one hand when playing or eating
- *Dragging one foot or walking on toes with one foot
- *Weakness on one side of the body

If you see any of these signs,
contact your doctor or your state's
early intervention program for assistance.

For more information: www.chasa.org

Pediatric Stroke can be frightening... Pediatric Stroke can be frightening...

1 in 4,000 babies will suffer a stroke before their first birthday, and many times before birth.

Some common signs of Pediatric Stroke:

- *Use of only one hand when playing or eating
- *Dragging one foot or walking on toes with one foot
- *Weakness on one side of the body

If you see any of these signs,
contact your doctor or your state's
early intervention program for assistance.

For more information: www.chasa.org

1 in 4,000 babies will suffer a stroke before their first birthday, and many times before birth.

Some common signs of Pediatric Stroke:

- *Use of only one hand when playing or eating
- *Dragging one foot or walking on toes with one foot
- *Weakness on one side of the body

If you see any of these signs,
contact your doctor or your state's
early intervention program for assistance.

For more information: www.chasa.org

Pediatric Stroke can be frightening... Pediatric Stroke can be frightening...

1 in 4,000 babies will suffer a stroke before their first birthday, and many times before birth.

Some common signs of Pediatric Stroke:

- *Use of only one hand when playing or eating
- *Dragging one foot or walking on toes with one foot
- *Weakness on one side of the body

If you see any of these signs,
contact your doctor or your state's
early intervention program for assistance.

For more information: www.chasa.org

1 in 4,000 babies will suffer a stroke before their first birthday, and many times before birth.

Some common signs of Pediatric Stroke:

- *Use of only one hand when playing or eating
- *Dragging one foot or walking on toes with one foot
- *Weakness on one side of the body

If you see any of these signs,
contact your doctor or your state's
early intervention program for assistance.

For more information: www.chasa.org