



Every bunny needs to know about...

## PEDIATRIC STROKE

**Infants have strokes, Kids have strokes, even unborn babies have strokes!**

### SOME COMMON SIGNS OF EARLY STROKE

- \*Dragging one foot or walking on toes with one foot.
- \*Weakness on one side of the body.
- \*Use of only one hand when playing or eating.

If you see any of these signs, contact your doctor or your state's early intervention program for assistance.

For more information: [www.chasa.org](http://www.chasa.org)



Every bunny needs to know about...

## PEDIATRIC STROKE

**Infants have strokes, Kids have strokes, even unborn babies have strokes!**

### SOME COMMON SIGNS OF EARLY STROKE

- \*Dragging one foot or walking on toes with one foot.
- \*Weakness on one side of the body.
- \*Use of only one hand when playing or eating.

If you see any of these signs, contact your doctor or your state's early intervention program for assistance.

For more information: [www.chasa.org](http://www.chasa.org)



Every bunny needs to know about...

## PEDIATRIC STROKE

**Infants have strokes, Kids have strokes, even unborn babies have strokes!**

### SOME COMMON SIGNS OF EARLY STROKE

- \*Dragging one foot or walking on toes with one foot.
- \*Weakness on one side of the body.
- \*Use of only one hand when playing or eating.

If you see any of these signs, contact your doctor or your state's early intervention program for assistance.

For more information: [www.chasa.org](http://www.chasa.org)



Every bunny needs to know about...

## PEDIATRIC STROKE


**Infants have strokes, Kids have strokes, even unborn babies have strokes!**

### SOME COMMON SIGNS OF EARLY STROKE

- \*Dragging one foot or walking on toes with one foot.
- \*Weakness on one side of the body.
- \*Use of only one hand when playing or eating.

If you see any of these signs, contact your doctor or your state's early intervention program for assistance.

For more information: [www.chasa.org](http://www.chasa.org)



Every bunny needs to know about...

## PEDIATRIC STROKE

**Infants have strokes, Kids have strokes, even unborn babies have strokes!**

### SOME COMMON SIGNS OF EARLY STROKE

- \*Dragging one foot or walking on toes with one foot.
- \*Weakness on one side of the body.
- \*Use of only one hand when playing or eating.

If you see any of these signs, contact your doctor or your state's early intervention program for assistance.

For more information: [www.chasa.org](http://www.chasa.org)



Every bunny needs to know about...

## PEDIATRIC STROKE

**Infants have strokes, Kids have strokes, even unborn babies have strokes!**

### SOME COMMON SIGNS OF EARLY STROKE

- \*Dragging one foot or walking on toes with one foot.
- \*Weakness on one side of the body.
- \*Use of only one hand when playing or eating.

If you see any of these signs, contact your doctor or your state's early intervention program for assistance.

For more information: [www.chasa.org](http://www.chasa.org)

