

## Spread Childhood Stroke Awareness...

1 in 2,800 babies will suffer a stroke before their first birthday, with some occurring before birth.

### Some common signs of early stroke:



- \*Use of only one hand when playing or eating
- \* Dragging one foot or walking on toes with one foot
- \* Weakness on one side of the body

If you see any of these signs, contact your doctor or your state's early intervention program for assistance.

For more information visit: [www.chasa.org](http://www.chasa.org)

## Spread Childhood Stroke Awareness...

1 in 2,800 babies will suffer a stroke before their first birthday, with some occurring before birth.

### Some common signs of early stroke:



- \*Use of only one hand when playing or eating
- \* Dragging one foot or walking on toes with one foot
- \* Weakness on one side of the body

If you see any of these signs, contact your doctor or your state's early intervention program for assistance.

For more information visit: [www.chasa.org](http://www.chasa.org)

## Spread Childhood Stroke Awareness...

1 in 2,800 babies will suffer a stroke before their first birthday, with some occurring before birth.

### Some common signs of early stroke:



- \*Use of only one hand when playing or eating
- \* Dragging one foot or walking on toes with one foot
- \* Weakness on one side of the body

If you see any of these signs, contact your doctor or your state's early intervention program for assistance.

For more information visit: [www.chasa.org](http://www.chasa.org)

## Spread Childhood Stroke Awareness...

1 in 2,800 babies will suffer a stroke before their first birthday, with some occurring before birth.

### Some common signs of early stroke:



- \*Use of only one hand when playing or eating
- \* Dragging one foot or walking on toes with one foot
- \* Weakness on one side of the body

If you see any of these signs, contact your doctor or your state's early intervention program for assistance.

For more information visit: [www.chasa.org](http://www.chasa.org)

## Spread Childhood Stroke Awareness...

1 in 2,800 babies will suffer a stroke before their first birthday, with some occurring before birth.

### Some common signs of early stroke:



- \*Use of only one hand when playing or eating
- \* Dragging one foot or walking on toes with one foot
- \* Weakness on one side of the body

If you see any of these signs, contact your doctor or your state's early intervention program for assistance.

For more information visit: [www.chasa.org](http://www.chasa.org)

## Spread Childhood Stroke Awareness...

1 in 2,800 babies will suffer a stroke before their first birthday, with some occurring before birth.

### Some common signs of early stroke:



- \*Use of only one hand when playing or eating
- \* Dragging one foot or walking on toes with one foot
- \* Weakness on one side of the body

If you see any of these signs, contact your doctor or your state's early intervention program for assistance.

For more information visit: [www.chasa.org](http://www.chasa.org)

## Spread Childhood Stroke Awareness...

1 in 2,800 babies will suffer a stroke before their first birthday, with some occurring before birth.

### Some common signs of early stroke:



- \*Use of only one hand when playing or eating
- \* Dragging one foot or walking on toes with one foot
- \* Weakness on one side of the body

If you see any of these signs, contact your doctor or your state's early intervention program for assistance.

For more information visit: [www.chasa.org](http://www.chasa.org)

## Spread Childhood Stroke Awareness...

1 in 2,800 babies will suffer a stroke before their first birthday, with some occurring before birth.

### Some common signs of early stroke:



- \*Use of only one hand when playing or eating
- \* Dragging one foot or walking on toes with one foot
- \* Weakness on one side of the body

If you see any of these signs, contact your doctor or your state's early intervention program for assistance.

For more information visit: [www.chasa.org](http://www.chasa.org)